

February 22, 2011

Multiple Choice	_____	(30 points)
Page 5	_____	(14 points)
Page 6	_____	(14 points)
Page 7	_____	(10 points)
Page 8	_____	(14 points)
Page 9	_____	(15 points)
Page 10	_____	(15 points)
Total	_____	(112 points)
Percent	_____	(100 %)

All work must be shown to receive credit. Give all answers to the correct number of significant figures

Avogadro's number = 6.022×10^{23} /mol
4 quarts = 1 gallon
36 in = 1 yard

Solubility Rules

- Alkali metals and NH_4^+ compounds are soluble.
- Nitrates(NO_3^-), acetates (CH_3CO_2^-), chlorates (ClO_3^-), perchlorates(ClO_4^-), and sulfates(SO_4^{2-}) are generally soluble (except for Sr^{2+} , Ca^{2+} , Ba^{2+} , Pb^{2+} , and Hg_2^{+2} sulfates).
- Chlorides(Cl^-), bromides(Br^-), iodides(I^-), are soluble (except for Silver(Ag^+), mercury(I)(Hg_2^{+2}), and lead(II)(Pb^{+2}) halides).
- Most compounds not included above are not soluble.
 - i.e. Sulfides(S^{2-}), carbonates(CO_3^{2-}), phosphates(PO_4^{3-}), chromates(CrO_4^{2-}), Oxides (O^{2-}), and Hydroxides(OH^-)
 - ($\text{Ca}(\text{OH})_2$, CaO , $\text{Sr}(\text{OH})_2$, SrO , $\text{Ba}(\text{OH})_2$ and BaO are slightly soluble.)

Grossmont College Periodic Table

VIIA NOBL
E
GASE
S

1 H 1.008	IIA																		1 H 1.008	2 He 4.002
3 Li 6.941	4 Be 9.012											5 B 10.81	6 C 12.01	7 N 14.01	8 O 16.00	9 F 19.00	10 Ne 20.18			
11 Na 23.00	12 Mg 24.30	IIIB	IVB	VB	VIB	VIIIB	VIII	VIII	VIII	IB	IIB	13 Al 27.00	14 Si 28.09	15 P 30.97	16 S 32.06	17 Cl 35.45	18 Ar 39.95			
19 K 39.10	20 Ca 40.08	21 Sc 44.96	22 Ti 47.90	23 V 50.94	24 Cr 52.00	25 Mn 54.94	26 Fe 55.85	27 Co 58.93	28 Ni 58.70	29 Cu 63.55	30 Zn 65.38	31 Ga 69.72	32 Ge 72.59	33 As 74.92	34 Se 78.96	35 Br 79.90	36 Kr 83.80			
37 Rb 85.47	38 Sr 87.62	39 Y 88.91	40 Zr 91.22	41 Nb 92.91	42 Mo 95.94	43 Tc (99)	44 Ru 101.1	45 Rh 102.9	46 Pd 106.4	47 Ag 107.9	48 Cd 112.4	49 In 114.8	50 Sn 118.7	51 Sb 121.8	52 Te 127.6	53 I 126.9	54 Xe 131.3			
55 Cs 132.9	56 Ba 137.3	57 La 138.9	72 Hf 178.5	73 Ta 180.9	74 W 183.9	75 Re 186.2	76 Os 190.2	77 Ir 192.2	78 Pt 195.1	79 Au 197.0	80 Hg 200.6	81 Tl 204.4	82 Pb 207.2	83 Bi 209.0	84 Po (209)	85 At (210)	86 Rn (222)			
87 Fr (223)	88 Ra 226.0	89 Ac 227.0	104 Rf (261)	105 Db (262)	106 Sg (263)	107 Bh (262)	108 Hs (265)	109 Mt (266)	110 ?? (269)											

Lanthanide series

58 Ce 140.1	59 Pr 140.9	60 Nd 144.2	61 Pm (147)	62 Sm 150.4	63 Eu 152.0	64 Gd 157.3	65 Tb 158.9	66 Dy 162.5	67 Ho 164.9	68 Er 167.3	69 Tm 168.9	70 Yb 173.0	71 Lu 175.0
90 Th 232.0	91 Pa 231.0	92 U 238.0	93 Np (237)	94 Pu (244)	95 Am (243)	96 Cm (247)	97 Bk (247)	98 Cf (251)	99 Es (252)	100 Fm (257)	101 Md (258)	102 No (259)	103 Lr (260)

Actinide series

Multiple Choice (30 points) – Give the best answer for each of the following questions.

- A consistent explanation of known observations is called
 - an experiment.
 - a prediction.
 - a hypothesis.
 - a theory.
- Which of the following statements does not describe a **chemical** property of oxygen?
 - The pressure is caused by collision of oxygen molecules with the sides of a container.
 - When coal is burned in oxygen, the process is called combustion.
 - Iron will rust in the presence of oxygen.
 - Oxygen combines with carbon to form carbon dioxide gas.
- A student measured the diameter of a sphere and determined the average value. His measurements are 6.17cm, 6.16cm, 6.16cm and 6.17cm. If the true diameter is 6.18 cm, what can be said about the student's results?
 - It is accurate and precise.
 - It is accurate but not precise.
 - It is precise but not accurate.
 - It is neither precise nor accurate.
- To the correct number of significant figures, what is the temperature reading on the following Celsius thermometer?
 - 15°C
 - 15.67°C
 - 16°C
 - 15.6°C
- Which of the following is **not** explained by Dalton's atomic theory?
 - the existence of more than one isotope of an element
 - the law of multiple proportions
 - conservation of mass during a chemical reaction
 - the law of definite proportions
- How many protons (p) and neutrons (n) are in an atom of calcium-46?
 - 26 p, 20 n
 - 46 p, 60 n
 - 20 p, 26 n
 - 20 p, 46 n
- In which set do all elements tend to form cations in binary ionic compounds?
 - Li, B, O
 - Mg, Cr, Pb
 - O, F, Cl
 - N, As, Bi



8. The solid compound, Na_2CO_3 , contains
- Na^+ , C^{4+} , and O^{2-} ions.
 - Na_2CO_3 molecules.
 - Na^+ ions and CO_3^{2-} ions.
 - Na_2^+ and CO_3^{2-} ions.
9. How many H^+ ions can the acid, H_2CO_3 , donate per molecule?
- 0
 - 1
 - 3
 - 2
10. Which one of the following statements about balanced equations is **false**? In a balanced reaction
- mass must be conserved.
 - net charge must be balanced on both sides of the reaction arrow.
 - molecules must be balanced on both sides of the reaction arrow.
 - atoms must be balanced on both sides of the reaction arrow.
11. Which statement about diluted solutions is **false**? When a solution is diluted
- the number of moles of solvent remains unchanged.
 - the concentration of the solution decreases.
 - the number of moles of solute remains unchanged.
 - the molarity of the solution decreases.
12. HBr , HCl , HClO_4 , KBr , and NaCl are all classified as
- strong electrolytes.
 - weak electrolytes.
 - acids.
 - nonelectrolytes.
13. What reagent could be used to separate Br^- from NO_3^- when added to an aqueous solution containing both?
- $\text{NaI} (aq)$
 - $\text{Ba}(\text{OH})_2 (aq)$
 - $\text{CuSO}_4 (aq)$
 - $\text{AgNO}_3 (aq)$
14. What is the oxidation number of the oxygen atom in H_2O_2 ?
- 1
 - +2
 - +1
 - 2
15. Which species functions as the oxidizing agent in the following reduction-oxidation reaction:
- $$5 \text{Fe}^{+2}(aq) + \text{MnO}_4^{-1}(aq) + 8 \text{H}^{+1}(aq) \rightarrow \text{Mn}^{+2}(aq) + 5 \text{Fe}^{+3}(aq) + 4 \text{H}_2\text{O}(aq)$$
- $\text{Mn}^{2+}(aq)$
 - $\text{H}^+(aq)$
 - $\text{Fe}^{2+}(aq)$
 - $\text{MnO}_4^-(aq)$

Problems

1. (5 points) Give the IUPAC name for the following compounds

- a. SrBr_2 strontium bromide
- b. AlPO_4 aluminum phosphate
- c. Cl_2O_7 dichlorine heptoxide
- d. LiClO_2 lithium chlorite
- e. $\text{V}(\text{NO}_3)_5$ vanadium(V) nitrate

2. (5 points) Write the correct formula for each of the following compounds

- a. Ammonium hypoiodite NH_4IO
- b. Zinc bromide ZnBr_2
- c. Ferric sulfate $\text{Fe}_2(\text{SO}_4)_3$
- d. Mercury(I) carbonate Hg_2CO_3
- e. Sulfur dioxide SO_2

3. (4 points) Perform the following calculation and report your answer with the correct number of significant figures.

$$\frac{6.34 + (90.3)(0.05442) + 943.8642}{(85.3992 - 86.000)} = \frac{6.34 + 4.91 + 943.8642}{85.3992 - 86.00} = \frac{955.12}{-0.601} = -1590$$

4. (8 points) Copper can be drawn into thin wires. How many meters of 34 gauge wire (diameter = 6.304×10^{-3} in) can be produced from the 8.01 lb of covallite, an ore of copper that is 66.0% copper by mass (Hint: Treat the wire as a cylinder: the density of copper is 8.95 g/cm^3 , figure out the mass of copper wire per unit length.)

$$\text{Volume of 1 in wire} = (\pi r^2)l = \left(\frac{\pi d^2}{4}\right)l = \left(\frac{\pi(6.304 \times 10^{-3} \text{ in})^2}{4}\right)1 \text{ in} = 3.121 \times 10^{-5} \text{ in}^3$$

$$\frac{\text{g Cu}}{\text{in}} = \frac{3.121 \times 10^{-5} \text{ in}^3}{\text{in wire}} \times \left(\frac{2.54 \text{ cm}}{1 \text{ in}}\right)^3 \times \frac{8.95 \text{ g}}{1 \text{ cm}^3} = \frac{4.58 \times 10^{-3} \text{ g Cu}}{\text{in wire}}$$

$$\begin{aligned} ? \text{ m wire} &= 8.01 \text{ lb ore} \times \frac{454 \text{ g ore}}{1 \text{ lb ore}} \times \frac{66.0 \text{ g Cu}}{100 \text{ g ore}} \times \frac{1 \text{ in wire}}{4.58 \times 10^{-3} \text{ g Cu}} \times \frac{2.54 \text{ cm wire}}{1 \text{ in wire}} \\ &\times \frac{1 \text{ m wire}}{100 \text{ cm wire}} \times \frac{1 \text{ km wire}}{1000 \text{ m wire}} = 13.3 \text{ km wire} \end{aligned}$$

5. (6 points) An element X forms both a dichloride (XCl_2) and a tetrachloride (XCl_4). Treatment of 10.00 g XCl_2 with excess chlorine forms 13.73 g XCl_4 . Calculate the atomic mass of X. Predict its identity.


$$13.73 \text{ g XCl}_4 - 10.00 \text{ g XCl}_2 = 3.73 \text{ g Cl added}$$

$$3.73 \text{ g Cl} \times \frac{1 \text{ mol Cl}}{35.45 \text{ g Cl}} \times \frac{1 \text{ mol XCl}_2}{2 \text{ mol Cl}} = 0.0526 \text{ mol XCl}_2$$

$$\begin{aligned} \text{molar mass XCl}_2 &= \frac{10.00 \text{ g XCl}_2}{0.0526 \text{ mol XCl}_2} = 190 \text{ gXCl}_2/\text{mol} \\ \text{molar mass} &= 190 \text{ amu} = 1 \text{ X}(\text{? amu/X}) + 2 \text{ Cl}(35.45 \text{ amu/Cl}) \end{aligned}$$

$$X = 118.7 \text{ probably tin}$$

6. (6 points) Complete the following double displacement reaction with balanced molecular, total ionic, and net ionic equations.




Balanced total ionic equation



Balanced net ionic equation




7. (4 points) Balance the following redox half reaction that occurs in basic solution



Is this an oxidation or a reduction?

reduction

8. (6 points) Balance the following redox reaction in acidic solution



1st half reaction



2nd half reaction



overall reaction in acid



9. (8 points) When 6.853 mg of a sex hormone containing C, H, and O was burned in a combustion analysis, 19.73 mg of CO₂ and 6.391 mg of H₂O were obtained. What is the empirical formula of the compound?


$$? \text{ mg C} = 19.73 \text{ mg CO}_2 \times \frac{1 \text{ mmol CO}_2}{44.01 \text{ mg CO}_2} \times \frac{1 \text{ mmol C}}{1 \text{ mmol CO}_2} = 0.4483 \text{ mmol C} \times \frac{12.01 \text{ mg C}}{1 \text{ mmol C}} = 5.384 \text{ mg C} \quad (78.56\% \text{ C}) \quad ($$

$$? \text{ mg H} = 6.391 \text{ mg H}_2\text{O} \times \frac{1 \text{ mmol H}_2\text{O}}{18.02 \text{ mg H}_2\text{O}} \times \frac{2 \text{ mmol H}}{1 \text{ mmol H}_2\text{O}} = 0.7093 \text{ mmol H} \times \frac{1.008 \text{ mg H}}{1 \text{ mmol H}} = 0.7150 \text{ mg H} \quad (10.43\% \text{ H})$$

$$? \text{ mg O} = (6.853 \text{ mg} - (5.384 \text{ mg C} + 0.7150 \text{ mg H})) = 6.853 \text{ mg} - 6.099 \text{ mg} = 0.754 \text{ mg O} \quad (11.00\% \text{ O})$$

$$? \text{ mmol O} = 0.754 \text{ mg O} \times \frac{1 \text{ mmol O}}{16.00 \text{ mg O}} = 0.0471 \text{ mmol O}$$

$$\frac{\text{C}_{0.4483} \text{H}_{0.7093} \text{O}_{0.0471}}{0.0471} \quad \frac{0.4483}{0.0471} \quad \frac{0.7093}{0.0471} \quad \frac{0.0471}{0.0471}$$



10. (5 points) How many grams of copper are in 50.0 mL of a 25.4% solution of copper (II) chloride with a density of 1.284 g/mL?

$$? g Cu = 50.0 mL soln \times \frac{1.284 g soln}{1 mL soln} \times \frac{25.4 g Cu}{100 g soln} = 16.3 g Cu$$

11. (10 points) Phencyclidine or angle dust has a molecular formula $C_{17}H_{25}N$. Answer the following questions regarding phencyclidine.

- a. Calculate the molar mass of phencyclidine.

$$\begin{aligned} \text{molar mass} &= 17 C(12.01 \text{ amu}/C) + 25 H(1.008 \text{ amu}/H) + 1 N(14.01 \text{ amu}/N) \\ &= 204.2 \text{ amu} + 25.20 \text{ amu} + 14.01 \text{ amu} = 243.4 \text{ amu} \end{aligned}$$

- b. Calculate the number of moles of carbon in 4.29 moles of phencyclidine.

$$? mol C = 4.29 mol phe \times \frac{17 mol C}{1 mol phe} = 72.9 mol C$$

- c. Calculate the number of molecules of phencyclidine that contains 675 atoms of hydrogen.

$$? molec phe = 675 atom H \times \frac{1 molec phe}{25 atom H} = 27 molec phe$$

- d. Calculate the mass of phencyclidine that contains 5.922×10^{24} atoms of carbon.


$$? g = 5.922 \times 10^{24} atom C \times \frac{1 mol C}{6.022 \times 10^{23} atom C} \times \frac{1 mol phe}{17 mol C} \times \frac{243.4 g phe}{1 mol phe} = 140.8 g phe$$

- e. Calculate the mass in grams of one molecule of phencyclidine.

$$? g phe = 1 molec phe \times \frac{1 mol phe}{6.022 \times 10^{23} molec phe} \times \frac{243.4 g phe}{1 mol phe} = 4.042 \times 10^{-22} g phe$$

12. (15 points) You mix 527.0 mL of 0.2754 M sodium carbonate with 400.0 mL of 0.6684 M chromium(III) chloride. Write the reaction and determine the number of grams of chromium(III) carbonate produced, and the final concentration of all ions in the solution.

Balanced chemical equation (Check with me before you go on to be sure this is correct.)



$$? \text{ mol Na}_2\text{CO}_3 = 527.0 \text{ mL} \times \frac{0.2754 \text{ mol Na}_2\text{CO}_3}{1000 \text{ mL}} = 0.1451 \text{ mol Na}_2\text{CO}_3$$

$$? \text{ mol CrCl}_3 = 400.0 \text{ mL} \times \frac{0.6684 \text{ mol CrCl}_3}{1000 \text{ mL}} = 0.2674 \text{ mol CrCl}_3$$

	X = 0.0484 mol		X=0.1337mol			
	3 Na ₂ CO ₃ (aq)	+	2 CrCl ₃ (aq)	→	6 NaCl(aq)	+ Cr ₂ (CO ₃) ₃ (s)
I	0.1451 mol		0.2674 mol		0 mol	0 mol
Δ	-3x		-2x		+6x	+ x
E	0.1451 - 3x		0.1671-2x		6x	1x
	=0.1451-3(.0484) =0 mol		=0.2674-2(.0484) =0.1707mol		=6(0.0484) =0.2904 mol	=0.0484 mol

$$0.0484 \text{ mol Cr}_2(\text{CO}_3)_3 \times \frac{284.0 \text{ g Cr}_2(\text{CO}_3)_3}{1 \text{ mol Cr}_2(\text{CO}_3)_3} = \boxed{13.7 \text{ g Cr}_2(\text{CO}_3)_3}$$

$$[\text{Na}^{+1}] = \frac{0.2904 \text{ mol Na}^{+1}}{0.9270 \text{ L solution}} = \boxed{0.3133 \text{ M Na}^{+1}}$$

$$[\text{CO}_3^{-2}] = \frac{0 \text{ mol CO}_3^{-2}}{0.9270 \text{ L solution}} = \boxed{0 \text{ M CO}_3^{-2}}$$

$$[\text{Cr}^{+3}] = \frac{0.1707 \text{ mol Cr}^{+3}}{0.9270 \text{ L solution}} = \boxed{0.1841 \text{ M Cr}^{+3}}$$

$$[\text{Cl}^{-1}] = \frac{3(0.1701 \text{ mol Cl}^{-1}) + 0.2904 \text{ mol Cl}^{-1}}{0.9270 \text{ L solution}} = \frac{0.8025 \text{ mol Cl}^{-1}}{0.9270 \text{ L solution}} = \boxed{0.8637 \text{ M Cl}^{-1}}$$

Moles Cr₂(CO₃)₃ produced 0.0484 mol Mass Cr₂(CO₃)₃ produced 13.7 g

Moles Na⁺¹ = 0.2904 mol

[Na⁺¹] = 0.3133M

Moles CO₃⁻² = 0 mol

[CO₃⁻²] = 0 M

Moles Cr⁺³ = 0.1707 mol

[Cr⁺³] = 0.1841 M

Moles Cl⁻¹ = 0.8025 mol

[Cl⁻¹] = 0.8637 M